MYCOTAXON

ISSN (print) 0093-4666 (online) 2154-8889 Mycotaxon, Ltd. ©2019

*January–March 2019—Volume 134, pp. 169–175*

https://doi.org/10.5248/134.169

# New records of *Hymenoscyphus, Parascutellinia,* and *Scutellinia* for Turkey

Ali Keleş*

*Department of Science and Mathematics Education, Yüzüncü Yıl University, Zeve Campus, Van, Turkey*

* Correspondence to: *alikeles61@yahoo.com*

Abstract—Three ascomycetous macrofungi—*Hymenoscyphus kathiae, Parascutellinia violacea,* and *Scutellinia crinita*—are recorded for the first time from Turkey, with *P. violacea* serving also the first record of the genus for the country. The species are described briefly and accompanied by photographs illustrating their macro- and micro-morphologies.

Key words—biodiversity, *Helotiaceae, Pyronemataceae,* Siirt, Van

## Introduction

Since studies on Turkish macromycota started in the second quarter of the 20th century, almost 2500 species have been reported. Considering the 15,000 macrofungi reported from Europe (Lukić 2009), it can easily be said that there is still much to be done.

This study aims to identify new records for the mycobiota of Turkey. During routine field studies some ascomycete samples were collected and identified as *Hymenoscyphus kathiae (Helotiaceae), Parascutellinia violacea (Pyronemataceae),* and *Scutellinia crinita (Pyronemataceae).* Among the 220 ascomycete species from Turkey covered by current checklists (Sesli & Denchev 2014, Solak & al. 2015) and recent publications on Turkish *Ascomycota* (Kaya & Uzun 2015, 2018; Acar & Uzun 2016; Kaya & al. 2016, 2018; Akçay & Uzun 2016; Doğan & al. 2016; Keleş & Oruç 2017; Uzun & al. 2017a,b, 2018a,b; Işık & Türkekul 2018), these three species are not cited and represent new records for the country.

## Materials & methods

Fungal specimens were collected from Baykan (Siirt) and from Gevaş and Gürpınar (Van) districts during 2012–2018. The fungi were photographed in their natural habitats and their ecological and morphological characteristics noted. Specimen samples were mounted in Congo red or Melzer's reagent and examined microscopically using a Leica DM500 light microscope. The fungi were identified after consulting Svrček (1975), Schumacher (1990), Korf (1999), Garcia & Van Vooren (2005), Medardi (2006), and Van Vooren & Hairaud (2009). The collections are kept in the fungarium of Yüzüncü Yıl University, Van, Turkey (VANF).

## Taxonomy

***Hymenoscyphus kathiae*** (Korf) Baral, Fl. Mediterr. 15: 66 (2005) Fig. 1

Apothecia 1–3 mm diam., subsessile to shortly stipitate, cupuliform, then more or less plane, margin smooth and glabrous. Hymenium smooth, pale yellow to golden yellow, outer surface almost concolorous with the hymenium or lighter. Stipe noticeably short, broad, yellowish white. Asci 90–110 × 7–10 µm, cylindrical, slightly attenuated at base and arising from croziers, generally 8-spored, some with fewer spores. Ascospores 9–14(–16) × 4–5 µm, ellipsoid, hyaline, smooth, some septate when mature, with two large or small guttules, uniseriate. Paraphyses cylindrical, rarely septate, hyaline, generally filled with yellowish guttules.

Specimen examined—**TURKEY, Siirt,** Baykan, central district, on wood remnants, 38°09'37"N 41°47'17"E, 769 m, 17.11.2012, A.K. 1877 (VANF).

Comments—*Hymenoscyphus kathiae* is macroscopically similar to *H. imberbis,* which differs in its ascospores with two large lipid droplets at the poles (Dimitrova & Baral 2005). In general, the morphological characteristics of our Turkish material agree with the descriptions by Dimitrova & Baral (2005) and Van Vooren & Hairaud (2009). This species is a new record for Turkey.

***Parascutellinia violacea*** (Velen.) Svrček, Česká Mykol. 29: 129 (1975) Fig. 2

Apothecia 3–6 mm diam., violet in colour, usually found in small groups; surface elliptical, smooth. Hymenium carmine red, with some violet tones. Asci 260–320 × 16–17 µm, cylindrical, operculate, inamyloid. Ascospores 24–29 × 12–14 µm, hyaline, elliptical, thick-walled, with one large guttule and two smaller guttules. Paraphyses multiseptate, branched, with expanded apex, containing a red-orange to slightly rosy pigment that turns greenish in Meltzer's. Paramarginal hairs yellowish-brown, fusiform, 75–150 µm, tightly crowded, fasciculate, multiseptate, with acute or obtuse apex, some well differentiated, others only poorly, with thickened walls ≤3 µm.




Fig 1. *Hymenoscyphus kathiae* (VANF A.K. 1877)
a. Ascomata; b. Ascospores; c. Tips of asci and paraphyses; d. Ascus;
e. Paraphyses; f. Ascus base.




Fig 2. *Parascutellinia violacea* (VANF A.K. 2961)
a. Ascomata; b. Ascospores; c. Paraphyses; d. Hairs; e. Asci

Specimen examined—**TURKEY, Van, Gürpınar,** Güzelsu village, river side, under *Salix* sp., 38°18'59"N–43°48'00"E, 1977 m, 05.10.2016, A.K. 2961 (VANF).

Comments— *Parascutellinia violacea* resembles *P. carneosanguinea,* which differs by carotenoid pigments within its paraphyses (Svrček 1975). Although

Van Vooren (2007) reported the same maximum sizes for the ascomata (5 mm) and asci as for *P. violacea,* some fruit bodies in our Turkish material measured almost 6 mm with asci up to 320 µm. Our specimens represent the first record of both genus and species for Turkey.




Fig 3. *Scutellinia crinita* (VANF A.K. 2963)
a. Ascomata; b. Ascospores; c. Ascal bases; d. Hairs; e. Asci;
f. Paraphyses; g. Apices of paraphyses.

***Scutellinia crinita*** (Bull.) Lambotte,
Mém. Soc. roy. Sci. Liège, Sér. 2, 14: 301 (1887) Fig. 3

Apothecia 3–7(–9) mm diam., disc-shaped, without stalk. Hymenium light orange, edges and outer surface covered with long dark brown hairs. Hairs 880–1550 × 22–40 µm, thick-walled, multiseptate, dark brownish, surface mostly smooth, sometimes cracked. Asci 207–250 × 12–22 µm, and cylindrical. Ascospores 16–22 × 15–18 µm, broadly rounded ellipsoid, slightly verrucose, containing many small droplets. Paraphyses filiform, clavate at apex, septate, somewhat branched.

Specimens examined—**TURKEY, Van, Gevaş,** at the vicinity of Kuzgunkiran tunnel, on rotten and damp wood under *Populus* sp., 38°22'09"N–42°47'40" E, 2052 m, 13.06.2018, A.K. 2963 (VANF).

Comments— The maximum size of ascomata for *Scutellinia crinita* was reported as 10 mm by Matočec & Antonić (1995) and 18 mm by Thompson (2013), but none of our ascomata measured >9 mm. This species represents a new record for Turkey.

**Acknowledgments**

The author would like to thank Prof. Dr. Ertuğrul Sesli, Prof. Dr. İbrahim Türkekul, and Dr. Shaun Pennycook for their helpful comments and careful review.

**Literature cited**

Acar İ, Uzun Y. 2016. *Peziza granularis* Donadini Türkiye Mikobiyotası için yeni bir kayıt. Yüzüncü Yıl Üniversitesi Fen Bilimleri Enstitüsü Dergisi 21 (1):39–42.

Akçay ME, Uzun Y. 2016. *Belonidium mollissimum (Lachnaceae):* Türkiye Mikotası için yeni bir tür. Mantar Dergisi 7(2): 118–121. https://doi.org/10.15318/Fungus.2016222679

Dimitrova EG, Baral HO. 2005. Checklist of Bulgarian *Helotiaceae (Ascomycetes).* Flora Mediterranea 15: 57–72.

Doğan HH, Bozok F, Taşkın H, Büyükalaca S 2016. Türkiye için beş yeni *Morchella* kaydı, Alatarım, 15 (1): 1–11.

Garcia G, Van Vooren N. 2005. Un discomycète inoperculé plutôt discret, *Pezoloma ciliifera*, et remarques sur le genre *Pezoloma.* Bulletin Mensuel de la Societe Linnéenne de Lyon 74: 115–129. https://doi.org/10.3406/linly.2005.13567

Işık H, Türkekul İ. 2018. A new record for Turkish mycota from Tokat Province: *Arachnopeziza aurelia* (Pers.) Fuckel. Journal of Fungus 9(1): 54–57.

Kaya A, Uzun Y. 2015. Six new genus records for Turkish *Pezizales* from Gaziantep Province. Turkish Journal of Botany 39: 506–511. https://doi.org/10.3906/bot-1409-3

Kaya A, Uzun Y. 2018. New contributions to the Turkish *Ascomycota*. Turkish Journal of Botany 42(5): 644–652. https://doi.org/10.3906/bot-1712-1

Kaya A, Uzun Y, Karacan Hİ, Yakar S. 2016. Contributions to Turkish *Pyronemataceae* from Gaziantep Province. Turkish Journal of Botany 40: 298–307. https://doi.org/10.3906/bot-1508-4

Kaya A, Uzun Y, Karacan İH, Yakar S. 2018. New additions to Turkish *Helotiales* and *Orbiliales*. Kastamonu University Journal of Forestry Faculty, 18(1): 46–52. https://doi.org/10.17475/kastorman.290359

Keleş A, Oruç Y. 2017. *Leucocoprinus brebissonii* (Godey) Locq, a new record for Turkish mycobiota. Anatolian Journal of Botany 1(2): 49–51. https://doi.org/10.30616/ajb.348736

Korf RP. 1999. *Pezoloma kathiae* sp. nov. (*Ascomycetes: Leotiales, Leotiaceae*), and its placement in a new subgenus, *Phaeopezoloma*. Mycotaxon, 73: 493–497.

Lukić N. 2009. The distribution and diversity of *Boletus* genus in Central Serbia. Kragujevac Journal of Science Sci 31: 59–68.

Matočec N, Antonić O. 1995. The genus *Scutellinia (Pezizales, Ascomycotina)* in Croatia: preliminary part. Natura Croatica, 4(1): 1–58.

Medardi G. 2006. Atlante fotografico degli ascomiceti d'Italia. Vicenza, Italy: Centro Studi Micologici.

Schumacher T. 1990. The genus *Scutellinia (Pyronemataceae).* Opera Botanica 101. 107 p.

Sesli E, Denchev CM 2014. Checklists of the myxomycetes, larger ascomycetes, and larger basidiomycetes in Turkey. 6th edn. Mycotaxon Checklists Online. (http://www.mycotaxon.com/resources/checklists/sesli-v106-checklist.pdf)

Solak MH, Işıloğlu M, Kalmış E, Allı H. 2015. Macrofungi of Turkey, checklist, vol. II. İzmir, Turkey: Üniversiteliler Ofset (in Turkish).

Svrček M. 1975. New or less known Discomycetes. II. Ceská Mykologie. 29(3):129–134.

Thompson PI. 2013. Ascomycetes in colour: found and photographed in mainland Britain. Indiana, USA: Xlibris.

Uzun Y, Acar İ, Akçay ME, Kaya A 2017a. Contributions to the macrofungi of Bingöl, Turkey. Turkish Journal of Botany 41: 516–534. https://doi.org/10.3906/bot-1611-14

Uzun Y, Karacan İH, Yakar S, Kaya A. 2017b. *Octospora* Hedw., a new genus record for Turkish *Pyronemataceae*. Anatolian Journal of Botany 1: 18–20. https://doi.org/10.30616/ajb.342114

Uzun Y, Karacan İH, Yakar S, Kaya A. 2018a. New bryophilic *Pyronemataceae* records for Turkish *Pezizales* from Gaziantep province. Anatolian Journal of Botany 2: 28–38. https://doi.org/10.30616/ajb.379549

Uzun Y, Yakar S, Karacan İH, Kaya A. 2018b. New additions to the Turkish *Pezizales.* Turkish Journal of Botany 42: 335–345.

Van Vooren N. 2007. Description et commentaire de quelques ascomycètes récoltés à la session de Lamoura (Jura). In Annales 2007. Compte rendu de la session mycologique de la FMBDS et des XXIe journées de la FAMM - Lamoura (Jura), septembre 2007 Publisher: Société Linnéenne de Lyon.

Van Vooren N, Hairaud M. 2009. Note sur *Hymenoscyphus kathiae* (*Helotiales*). Ascomycete.org 1(4): 11–15.